arXiv:1403.6737v2 [math.FA] 28 Mar 2014

On a conjecture of Pisier on the analyticity of semigroups

Cédric Arhancet

Abstract

We show that the analyticity of semigroups $(T_t)_{t\geqslant 0}$ of selfadjoint contractive Fourier multipliers on L^p-spaces of compact abelian groups is preserved by the tensorisation of the identity operator of a Banach space for a large class of K-convex Banach spaces, answering partially a conjecture of Pisier. We also give versions of this result for some semigroups of Schur multipliers and Fourier multipliers on noncommutative L^p-spaces. Finally, we give a precise description of semigroups of Schur multipliers to which the result of this paper can be applied.

1 Introduction

In the early eighties, in a famous paper on the geometry of Banach spaces, Pisier [Pis3] showed that a Banach space X does not contain ℓ_n^1's uniformly if and only if the tensorisation $P\otimes Id_X$ of the Rademacher projection

$$\begin{array}{rccc} P\colon & L^2(\Omega) & \longrightarrow & L^2(\Omega) \\ & f & \longmapsto & \displaystyle\sum_{k=1}^{\infty}\Big(\int_\Omega f\varepsilon_k\Big)\varepsilon_k \end{array}$$

induces a bounded operator on the Bochner space $L^2(\Omega,X)$ where Ω is a probability space and where $\varepsilon_1,\varepsilon_2,\dots$ is a sequence of independant random variables with $P(\varepsilon_k=1)=P(\varepsilon_k=-1)=\frac{1}{2}$. Such a Banach space X is called K-convex. The heart of his proof relies on the fact, proved by himself in his article, that if X is a K-convex Banach space then any w^*-continuous semigroup $(T_t)_{t\geqslant 0}$ of positive unital selfadjoint Fourier multipliers on a locally compact abelian group G induces a strongly continuous bounded analytic semigroup $(T_t\otimes Id_X)_{t\geqslant 0}$ of contractions on the Bochner space $L^p(G,X)$ where $1<p<\infty$. In 1981, in the seminars [Pis1] and [Pis2] which announced the results of his paper, he stated several natural questions raised by his work. In particular, he conjectured that the same property holds for any w^*-continuous semigroup $(T_t)_{t\geqslant 0}$ of selfadjoint contractive operators on a space $L^\infty(\Omega)$ where Ω is a measure space (see also the recent preprint [Xu, Problem 11] for a more general question). Note that it is well known that such a semigroup induces a strongly continuous bounded analytic semigroup of contractions on the associated L^p-space $L^p(\Omega)$ and the conjecture says that the property of analyticity is preserved by the tensorisation of the identity Id_X of a K-convex Banach space X.

The author is supported by the research program ANR 2011 BS01 008 01.
2010 *Mathematics subject classification:* 46L51, 46L07, 47D03.
Key words: noncommutative L^p-spaces, operator spaces, analytic semigroups, K-convexity, Fourier multipliers, Schur multipliers.

Using operator space theory (see [ER], [Pau] and [Pis7]), a quantised theory of Banach spaces, we are able to give the following partial answer to this purely Banach spaces question. First of all, let us recall that an operator space E is OK-convex if the vector valued Schatten space $S^p(E)$ is K-convex for some (equivalently all) $1 < p < \infty$. It means that the Rademacher projection P is completely bounded. This notion was introduced by [JuP] and is the noncommutative version of the property of K-convexity. Our main result is the following theorem.

Theorem 1.1 *Suppose that G is a compact abelian group. Let $(T_t)_{t\geqslant 0}$ be a w^*-continuous semigroup of selfadjoint contractive Fourier multipliers on $L^\infty(G)$. Let X be a K-convex Banach space isomorphic to a Banach space E which admits an OK-convex operator space structure. Consider $1 < p < \infty$. Then $(T_t)_{t\geqslant 0}$ induces a strongly continuous bounded analytic semigroup $(T_t \otimes Id_X)_{t\geqslant 0}$ of contractions on the Bochner space $L^p(G, X)$.*

This result can be used, by example, in the case where the Banach space X is an L^q-space or a Schatten space S^q with $1 < q < \infty$. Our methods also give a result for some w^*-continuous semigroups of Schur multipliers and a generalization for semigroups of Fourier multipliers on amenable discrete groups.

The paper is organized as follows. Section 2 gives a brief presentation of vector valued noncommutative L^p-spaces, Fourier multipliers on group von Neumann algebras and Schur multipliers. We introduce here some notions which are relevant to our paper. The next section 3 contains a proof of Theorem 1.1. Finally, in Section 4, we describe the semigroups of Schur multipliers to which the results of this paper can be applied. This result is of independent interest.

2 Preliminaries

The readers are referred to [ER], [Pau] and [Pis7] for details on operator spaces and completely bounded maps and to the survey [PX] for noncommutative L^p-spaces and the references therein.

If $T\colon E \to F$ is a completely bounded map between two operators spaces E and F, we denote by $\|T\|_{cb,E\to F}$ its completely bounded norm.

The theory of vector valued noncommutative L^p-spaces was initiated by Pisier [Pis5] for the case where the underlying von Neumann algebra is hyperfinite and equipped with a faithful normal semifinite trace. Suppose $1 \leqslant p < \infty$. Under theses assumptions, for any operator space E, we can define by complex interpolation

$$L^p(M, E) = \big(M \otimes_{\min} E, L^1(M)\widehat{\otimes}E\big)_{\frac{1}{p}}, \tag{2.1}$$

where $\otimes_{\min}$ and $\widehat{\otimes}$ denote the injective and the projective tensor product of operator spaces.

If $M = B(\ell_I^2)$ is the algebra of bounded operators on the Hilbert space ℓ_I^2 equipped with its canonical trace, we obtain the vector valued Schatten space $S_I^p(E) = L^p(B(\ell_I^2), E)$. Sometimes, we will use the notation $S^p(B(\ell_I^2)^{\otimes n}, E)$ for the space $L^p(B(\ell_I^2)^{\otimes n}, E)$.

Note the following vector valued extension property of completely positive maps between noncommutative L^p-spaces, see [Pis4].

Proposition 2.1 *Suppose $1 < p < \infty$. Let M and N be hyperfinite von Neumann algebras equipped with normal faithful semifinite traces and let E be an operator space. Let $T\colon M \to N$ be a trace preserving unital normal completely positive map. Then the operator $T \otimes Id_E$ extends to a bounded operator from $L^p(M, E)$ into $L^p(N, E)$ and we have*

$$\|T \otimes Id_E\|_{L^p(M,E)\to L^p(N,E)} \leqslant \|T\|_{L^p(M)\to L^p(N)}.$$

In particular, this result applies to canonical conditional expectations. Recall that a linear map $T\colon L^p(M) \to L^p(M)$ defined on a noncommutative L^p-space where $1 < p < \infty$ is said to be regular [Pis4] if for any operator space E the linear map $T \otimes Id_E\colon L^p(M,E) \to L^p(M,E)$ is bounded. The above proposition give an example of regular map.

Suppose that G is a discrete group. We denote by e_G the neutral element of G. For $f \in \ell^1(G)$, we write L_f for the left convolution by f acting on $\ell^2(G)$ by:

$$L_f(h)(g') = \sum_{g\in G} f(g)h(g^{-1}g')$$

for any $h \in \ell^2(G)$ and any $g' \in G$. Let $\mathrm{VN}(G)$ be the von Neumann algebra generated by $\{L_f\}_{f\in\ell^1(G)}$. This $\mathrm{VN}(G)$ is called the group von Neumann algebra of G and is equal to the von Neumann algebra generated by $\{\lambda_g\}_{g\in G}$, where λ_g is the left translation acting on $\ell^2(G)$ defined by $\lambda_g(h)(g') = h(g^{-1}g')$. It is an finite algebra with normalized trace given by

$$\tau_G(x) = \left\langle \epsilon_{e_G}, x(\epsilon_{e_G})\right\rangle_{\ell^2_G}$$

where $(\epsilon_g)_{g\in G}$ is the canonical basis of ℓ^2_G and $x \in VN(G)$.

For a compact abelian group G, the Fourier transform of $f \in \ell^1(G)$ is defined on the dual group $\hat{G}$ of G by

$$\hat{f}(\gamma) = \int_G f(x)\overline{\gamma(x)}dx$$

for $\gamma \in \hat{G}$. In this case, $(\mathrm{VN}(G),\, \tau_G)$ is equivalent as a von Neumann algebra to $L^\infty(\hat{G})$ with the usual integration on the dual group $\hat{G}$ of G under the mapping

$$\begin{array}{ccc} \mathrm{VN}(G) & \longrightarrow & L^\infty(\hat{G}) \\ L_f & \longmapsto & \hat{f}. \end{array} \tag{2.2}$$

Moreover, it is well known that G is discrete if and only if the dual group $\hat{G}$ is compact.

Let G be a discrete group. Recall that the von Neumann algebra $VN(G)$ is hyperfinite if and only if G is amenable [SS, Theorem 3.8.2]. A Fourier multiplier is a normal linear map $T\colon VN(G) \to VN(G)$ such that there exists a complex function $\varphi\colon G \to \mathbb{C}$ such that for any $g \in G$ we have $T(\lambda_g) = \varphi_g\lambda_g$. In this case, we denote T by

$$M_\varphi :\ \begin{array}{ccc} VN(G) & \longrightarrow & VN(G) \\ \lambda_g & \longmapsto & \varphi_g\lambda_g. \end{array}$$

By [DCH, Corollary 1.8], if the discrete group G is amenable, every contractive Fourier multiplier $M_\varphi\colon \mathrm{VN}(G) \to VN(G)$ is completely contractive.

If I is an index set and if E is a vector space, we write $\mathbb{M}_I$ for the space of the $I \times I$ matrices with entries in $\mathbb{C}$ and $\mathbb{M}_I(E)$ for the space of the $I \times I$ matrices with entries in E.

Let $A = [a_{ij}]_{i,j\in I}$ be a matrix of $\mathbb{M}_I$. By definition, the Schur multiplier on $B\big(\ell^2_I\big)$ associated with this matrix is the unbounded linear operator M_A whose domain is the space of all $B = [b_{ij}]_{i,j\in I}$ of $B\big(\ell^2_I\big)$ such that $[a_{ij}b_{ij}]_{i,j\in I}$ belongs to $B\big(\ell^2_I\big)$, and whose action on $B = [b_{ij}]_{i,j\in I}$ is given by $M_A(B) = [a_{ij}b_{ij}]_{i,j\in I}$. For all $i, j \in I$, the matrix e_{ij} belongs to $D(M_A)$, hence M_A is densely defined for the weak* topology. Suppose $1 \leqslant p < \infty$. If for any $B \in S^p_I$, we have $B \in D(M_A)$ and the matrix $M_A(B)$ represents an element of S^p_I, by the closed graph theorem, the matrix A of $\mathbb{M}_I$ defines a bounded Schur multiplier $M_A\colon S^p_I \to S^p_I$. We have a similar statement for bounded Schur multipliers on $B\big(\ell^2_I\big)$. Recall that every contractive Schur

multiplier $M_A\colon B(\ell_I^2) \to B(\ell_I^2)$ is completely contractive (see [Pau]). We say that a matrix A of $\mathbb{M}_I$ induces a completely positive Schur multiplier $M_A\colon B(\ell_I^2) \to B(\ell_I^2)$ if and only if for any finite set $F \subset I$ the matrix $[a_{i,j}]_{i,j\in F}$ is positive (see [Pau]).

Let M be a von Neumann algebra equipped with a semifinite normal faithful trace τ. Suppose that $T\colon M \to M$ is a normal contraction. We say that T is selfadjoint if for all $x, y \in M \cap L^1(M)$ we have

$$\tau\big(T(x)y^*\big) = \tau\big(x(T(y))^*\big).$$

In this case, it is not hard to show that the restriction $T|M \cap L^1(M)$ extends to a contraction $T\colon L^1(M) \to L^1(M)$. By complex interpolation, for any $1 \leqslant p \leqslant \infty$, we obtain a contractive map $T\colon L^p(M) \to L^p(M)$. Moreover, the operator $T\colon L^2(M) \to L^2(M)$ is selfadjoint. If $T\colon M \to M$ is actually a normal selfadjoint complete contraction, it is easy to see that the map $T\colon L^p(M) \to L^p(M)$ is completely contractive for any $1 \leqslant p \leqslant \infty$. It is not difficult to show that a contractive Fourier multiplier $M_\varphi\colon VN(G) \to VN(G)$ is selfadjoint if and only if $\varphi\colon G \to \mathbb{C}$ is a real function. Finally, one can prove that a contractive Schur multiplier M_φ is selfadjoint if and only if φ is a real function.

3 Analyticity of semigroups on vector valued L^p-spaces

Let X be a Banach space. A strongly continuous semigroup $(T_t)_{t\geqslant 0}$ is called bounded analytic if there exist $0 < \theta < \frac{\pi}{2}$ and a bounded holomorphic extension

$$\begin{array}{ccc} \Sigma_\theta & \longrightarrow & B(X) \\ z & \longmapsto & T_z \end{array}$$

where $\Sigma_\theta = \{z \in \mathbb{C}^* \ : \ |\mathrm{Arg}(z)| < \theta\}$ denotes the open sector of angle 2θ around the positive real axis $\mathbb{R}_+$. See [EN] and [Haa] for more information on this notion. We need the following theorem which is a corollary [Pis3, Lemma 4] of a result of Beurling [Beu] (see also [Fac] and [Hin]).

Theorem 3.1 *Let X be a Banach space. Let $(T_t)_{t\geqslant 0}$ be a strongly continuous semigroup of contractions on X. Suppose that there exists some integer $n \geqslant 1$ such that for any $t > 0$*

$$\big\|(Id_X - T_t)^n\big\|_{X\to X} < 2^n.$$

Then the semigroup $(T_t)_{t\geqslant 0}$ is bounded analytic.

Moreover, we recall the following lemma, see [DJT, Lemma 13.12] and [Pis3, Lemma 1.5].

Lemma 3.2 *Suppose that X is a K-convex Banach space. Then there exist a real number $0 < \rho < 2$ and an integer $n \geqslant 1$ such that if $P_1, \ldots, P_n$ is any finite collection of mutually commuting norm one projections on X, then*

$$\left\| \prod_{1\leqslant k\leqslant n} (Id_X - P_k) \right\|_{X\to X} \leqslant \rho^n.$$

We will use the useful next 'absorption Lemma' which is a variant of [Arh2, Proposition 3.4]. The proof is left to the reader.

Lemma 3.3 *Suppose $1 \leqslant p < \infty$. Let E be an operator space. For any positive integer $n \geqslant 1$ and any matrix $x \in \mathbb{M}_I(E)$ finitely supported on $I \times I$, we have*

$$\left\| \sum_{i,j\in I} e_{ij} \otimes e_{ij} \otimes \cdots \otimes e_{ij} \otimes x_{ij} \right\|_{S^p(B(\ell_I^2)^{\otimes n},E)} = \left\| \sum_{i,j\in I} e_{ij} \otimes x_{ij} \right\|_{S_I^p(E)}. \tag{3.1}$$

Moreover, for any regular Schur multiplier $M_A\colon S_I^p \to S_I^p$ and any positive integer $n \geqslant 1$ we have

$$\left\|M_A^n \otimes Id_E\right\|_{S_I^p(E)\to S_I^p(E)} \leqslant \left\|M_A^{\otimes n} \otimes Id_E\right\|_{S^p(B(\ell_I^2)^{\otimes n},E)\to S^p(B(\ell_I^2)^{\otimes n},E)}. \tag{3.2}$$

We also need the following transfer results of Neuwirth and Ricard [NR] between Fourier multipliers and Schur multipliers. Let G be a discrete amenable group. If $\varphi\colon G \to \mathbb{C}$ is a complex function, we denote by $\check{M}_\varphi$ the Schur multiplier defined by the symbol $\check{\varphi} \in \ell^\infty(G \times G)$ defined by $\check{\varphi}(g,h) = \varphi(gh^{-1})$ where $g, h \in G$. Then, for any $1 \leqslant p \leqslant \infty$, we have

$$\left\|M_\varphi \otimes Id_E\right\|_{L^p(\mathrm{VN}(G),E)\to L^p(\mathrm{VN}(G),E)} \leqslant \left\|\check{M}_\varphi \otimes Id_E\right\|_{S_G^p(E)\to S_G^p(E)} \tag{3.3}$$

and

$$\left\|M_\varphi \otimes Id_E\right\|_{cb,L^p(\mathrm{VN}(G),E)\to L^p(\mathrm{VN}(G),E)} = \left\|\check{M}_\varphi \otimes Id_E\right\|_{cb,S_G^p(E)\to S_G^p(E)}. \tag{3.4}$$

Using the identification (2.2), we see that Theorem 1.1 is a particular case of the following more general result, which improves a part of [Arh2, Theorem 5.1].

Theorem 3.4 *Suppose that G is an amenable discrete group. Let $(T_t)_{t\geqslant 0}$ be a w^*-continuous semigroup of selfadjoint contractive Fourier multipliers on the group von Neumann algebra $\mathrm{VN}(G)$. Suppose that E is an OK-convex operator space. Consider $1 < p < \infty$. Then $(T_t)_{t\geqslant 0}$ induces a strongly continuous bounded analytic semigroup $(T_t \otimes Id_E)_{t\geqslant 0}$ of contractions on the noncommutative vector valued L^p-space $L^p(\mathrm{VN}(G), E)$.*

Proof : We consider the associated semigroup $\left(\check{T}_t\right)_{t\geqslant 0}$ of selfadjoint contractive Schur multipliers on the space $B(\ell_G^2)$. Using (3.4), we see that

$$\left\|\check{T}_t\right\|_{S_G^p(E)\to S_G^p(E)} \leqslant \left\|\check{T}_t\right\|_{cb,S_G^p(E)\to S_G^p(E)} = \left\|T_t\right\|_{cb,L^p(\mathrm{VN}(G),E)\to L^p(\mathrm{VN}(G),E)} \leqslant 1.$$

In the sequel, we denote by $(\check{T}_t)^\circ$ the Schur multiplier defined by the adjoint matrix of the matrix of the Schur multiplier $\check{T}_t$. As the proof of [Arh1, Corollary 4.3], for any $t \geqslant 0$, there exists Schur multipliers $S_{1,t}$ and $S_{2,t}$ on $B(\ell_G^2)$ such that

$$W_t = \begin{bmatrix} S_{1,t} & \check{T}_t \\ (\check{T}_t)^\circ & S_{2,t} \end{bmatrix},$$

is a completely positive unital self-adjoint Schur multiplier on $B\big(\ell^2_{\{1,2\}\times G}\big)$. Note that, for any $t \geqslant 0$, we have

$$\left(W_{\frac{t}{2}}\right)^2 = \begin{bmatrix} S_{1,\frac{t}{2}} & \check{T}_{\frac{t}{2}} \\ \left(\check{T}_{\frac{t}{2}}\right)^\circ & S_{2,\frac{t}{2}} \end{bmatrix}^2 = \begin{bmatrix} S_{1,\frac{t}{2}}^2 & (\check{T}_{\frac{t}{2}})^2 \\ \left(\check{T}_{\frac{t}{2}}\right)^{\circ 2} & S_{2,\frac{t}{2}}^2 \end{bmatrix} = \begin{bmatrix} S_{1,\frac{t}{2}}^2 & \check{T}_t \\ (\check{T}_t)^\circ & S_{2,\frac{t}{2}}^2 \end{bmatrix}. \tag{3.5}$$

By [Ric] (see also [JMX, Section 10.B] for some related information), for any $t \geqslant 0$, the Schur multiplier $(W_{\frac{t}{2}})^2$ admits a Rota dilation

$$\big((W_{\frac{t}{2}})^2\big)^k = Q\mathbb{E}_k\pi, \qquad k \geqslant 1$$

where $\pi\colon M_2(B(\ell_G^2)) \to M$ is a normal unital faithful $*$-representation into a hyperfinite von Neumann algebra (equipped with a normalized trace) which preserve the traces, where $Q\colon M \to M_2(B(\ell_G^2))$ is the conditional expectation associated with π and where the $\mathbb{E}_k$'s are conditional expectations onto von Neumann subalgebras of M. In particular, we have

$$(W_{\frac{t}{2}})^2 = Q\mathbb{E}_1\pi.$$

We infer that

$$Id_{M_2(B(\ell_G^2))} - \big(W_{\frac{t}{2}}\big)^2 = Q(Id_M - \mathbb{E}_1)\pi.$$

Let n be a positive integer. We deduce that

$$\Big(Id_{M_2(B(\ell_G^2))} - \big(W_{\frac{t}{2}}\big)^2\Big)^{\otimes n} = Q(Id_M - \mathbb{E}_1)^{\otimes n}\pi. \tag{3.6}$$

For any integer $1 \leqslant k \leqslant n$, we let

$$\Pi_k = Id_{L^p(M)} \otimes \cdots \otimes Id_{L^p(M)} \otimes \mathbb{E}_1 \otimes Id_{L^p(M)} \otimes \cdots \otimes Id_{L^p(M)}.$$

By Proposition 2.1, we deduce that the $\Pi_k \otimes Id_E$'s are well-defined and form a family of mutually commuting contractive projections on the Banach space $L^p(M^{\otimes n}, E)$. Moreover, by [Arh2, Proposition 3.5], the latter space is K-convex. Now, we choose the integer $n \geqslant 1$ and $0 < \rho < 2$ as in Lemma 3.2. Hence, we obtain that

$$\Big\| \prod_{1\leqslant k\leqslant n} \Big(Id_{L^p(M^{\otimes n},E)} - (\Pi_k \otimes Id_E)\Big)\Big\|_{L^p(M^{\otimes n},E)\to L^p(M^{\otimes n},E)} \leqslant \rho^n. \tag{3.7}$$

Furthermore, it is easy to see that

$$\big(Id_{L^p(M)} - \mathbb{E}_1\big)^{\otimes n} \otimes Id_E = \prod_{1\leqslant k\leqslant n} \Big(Id_{L^p(M^{\otimes n},E)} - (\Pi_k \otimes Id_E)\Big).$$

Now, using (3.5), (3.2), (3.6) and (3.7) we obtain that

$$\begin{aligned}
\Big\|\big(Id_{S_G^p} - \check{T}_t\big)^n \otimes Id_E\Big\|_{S_G^p(E)\to S_G^p(E)} &\leqslant \Big\|\Big(Id_{S_2^p(S_G^p)} - \big(W_{\frac{t}{2}}\big)^2\Big)^n \otimes Id_E\Big\|_{S_2^p(S_G^p(E))\to S_2^p(S_G^p(E))} \\
&\leqslant \Big\|\Big(Id_{S_2^p(S_G^p)} - \big(W_{\frac{t}{2}}\big)^2\Big)^{\otimes n} \otimes Id_E\Big\|_{S^p(B(\ell^2_{\{1,2\}\times G})^{\otimes n},E)\to S^p(B(\ell^2_{\{1,2\}\times G})^{\otimes n},E)} \\
&= \Big\|Q(Id_{L^p(M)} - \mathbb{E}_1)^{\otimes n}\pi \otimes Id_E\Big\|_{S^p(B(\ell^2_{\{1,2\}\times G})^{\otimes n},E)\to S^p(B(\ell^2_{\{1,2\}\times G})^{\otimes n},E)} \\
&\leqslant \Big\|\big(Id_{L^p(M)} - \mathbb{E}_1\big)^{\otimes n} \otimes Id_E\Big\|_{L^p(M^{\otimes n},E)\to L^p(M^{\otimes n},E)} \\
&= \Big\| \prod_{1\leqslant k\leqslant n} \Big(Id_{L^p(M^{\otimes n},E)} - (\Pi_k \otimes Id_E)\Big)\Big\|_{L^p(M^{\otimes n},E)\to L^p(M^{\otimes n},E)} \leqslant \rho^n
\end{aligned}$$

Hence, using (3.3), for any $t \geqslant 0$, we obtain

$$\Big\| \big(Id_{S^p_G} - T_t\big)^n \otimes Id_E \Big\|_{L^p(\mathrm{VN}(G),E)\to L^p(\mathrm{VN}(G),E)} \leqslant \Big\| \big(Id_{S^p_G} - \check{T}_t\big)^n \otimes Id_E \Big\|_{S^p_G(E)\to S^p_G(E)} \leqslant \rho^n.$$

We conclude by Beurling Theorem 3.1. ∎

The next result is an improvement of [Arh2, Theorem 3.7] and can be proved as Theorem 3.4.

Theorem 3.5 *Let $(T_t)_{t\geqslant 0}$ be a w^*-continuous semigroup of selfadjoint contractive Schur multipliers on $B(\ell^2_I)$. Suppose that E is an OK-convex operator space. Consider $1 < p < \infty$. Then $(T_t)_{t\geqslant 0}$ induces a strongly continuous bounded analytic semigroup $(T_t \otimes Id_E)_{t\geqslant 0}$ of contractions on the vector valued Schatten space $S^p_I(E)$.*

Remark 3.6 *Wo does not know if any K-convex Banach space X is isomorphic to a Banach space E admitting an operator space structure such that the Banach space $S^p(E)$ is K-convex for $1 < p < \infty$ (i.e. E is OK-convex). Moreover, it would be also interesting to examine a similar question for other Banach spaces properties (UMD, cotype...).*

4 Semigroups of contractive selfadjoint Schur multipliers

The description of self-adjoint contractive Schur multipliers on $B(\ell^2_I)$ (and more generally of contractive Schur multipliers) is well-known and essentially goes back to Grothendieck and was rediscovered by many authors, see [Pis6, Chapter 5] for more information. Here, we give a continuous version of this result which precisely describes the semigroups of Schur multipliers of Theorem 3.5.

Theorem 4.1 *Suppose that A is a matrix of $\mathbb{M}_I$. For any $t \geqslant 0$, let T_t be the unbounded Schur multipliers on $B\big(\ell^2_I\big)$ associated with the matrix*

$$\Big[e^{-ta_{ij}}\Big]_{i,j\in I}. \tag{4.1}$$

The semigroup $(T_t)_{t\geqslant 0}$ extends to a semigroup of selfadjoint contractive Schur multipliers $T_t\colon B\big(\ell^2_I\big) \to B\big(\ell^2_I\big)$ if and only if there exists a Hilbert space H and two families $(\alpha_i)_{i\in I}$ and $(\beta_j)_{j\in I}$ of elements of H such that $a_{ij} = \|\alpha_i - \beta_j\|^2_H$ for any $i, j \in I$.

In this case, the Hilbert space may be chosen as a real Hilbert space and moreover, $(T_t)_{t\geqslant 0}$ is a w^*-continuous semigroup.

Proof : First, suppose that the semigroup $(T_t)_{t\geqslant 0}$ extends to a semigroup of selfadjoint contractive Schur multipliers $T_t\colon B\big(\ell^2_I\big) \to B\big(\ell^2_I\big)$. In particular, for any integer $n \geqslant 1$, the Schur multiplier $T_{\frac{1}{n}}\colon B\big(\ell^2_I\big) \to B\big(\ell^2_I\big)$ is contractive and selfadjoint. Thus, as the proof of [Arh2, Corollary 4.3], we can find matrices $S_{1,n}$, $S_{2,n} \in \mathbb{M}_I$ such that the block matrix

$$\begin{bmatrix} S_{1,n} & \big[e^{-\frac{a_{ij}}{n}}\big] \\ \big[e^{-\frac{a_{ji}}{n}}\big] & S_{2,n} \end{bmatrix}$$

defines a unital selfadjoint completely positive Schur multiplier on $B\big(\ell^2_{\{1,2\}\times I}\big)$. This matrix identifies to a matrix $[b_{n,k,i,m,j}]_{(k,i)\in\{1,2\}\times I,(m,j)\in\{1,2\}\times I} \in \mathbb{M}_{\{1,2\}\times I}$ such that for any $i, j \in I$

$$b_{n,1,i,1,j} = (S_{1,n})_{i,j}, \qquad b_{n,1,i,2,j} = e^{-\frac{a_{ij}}{n}},$$
$$b_{n,2,i,1,j} = e^{-\frac{a_{ji}}{n}}, \qquad b_{n,2,i,2,j} = (S_{2,n})_{i,j}.$$

Then it is obvious that the map

$$\begin{array}{ccc} (\{1,2\}\times I)\times(\{1,2\}\times I) & \longrightarrow & \mathbb{R} \\ \big((k,i),(m,j)\big) & \longmapsto & n(1-b_{n,k,i,m,j}) \end{array}$$

is a real-valued negative definite kernel which vanishes on the diagonal of $(\{1,2\}\times I)\times(\{1,2\}\times I)$. By [BCR, Proposition 3.2, Chapter 3], there exist a real Hilbert space H_n and a map $\xi^n : \{1,2\}\times I \to H_n$ such that

$$\big\|\xi^n_{k,i}-\xi^n_{m,j}\big\|^2_{H_n} = n\big(1-b_{n,k,i,m,j}\big), \quad k,m\in\{1,2\},\ i,j\in I.$$

We can assume that $\xi^n_{2,1}=0$ for any integer $n\geqslant 1$. For any $a\in\mathbb{R}$, we have $\frac{1-e^{-ta}}{t}\xrightarrow[t\to 0]{} a$. Hence, for any $i,j\in I$, we see that

$$\big\|\xi^n_{1,i}-\xi^n_{2,j}\big\|^2_{H_n} = n\big(1-b_{n,1,i,2,j}\big) = n\big(1-e^{-\frac{a_{ij}}{n}}\big)\xrightarrow[n\to\infty]{} a_{ij}.$$

Since $\xi^n_{2,1}=0$, we infer that $\big(\|\xi^n_{1,i}\|_{H_n}\big)_{n\geqslant 1}$ is a bounded sequence for each $i\in I$. Note that for any $i,j\in I$ we have

$$\big\|\xi^n_{2,j}\big\|_{H_n} \leqslant \big\|\xi^n_{1,i}-\xi^n_{2,j}\big\|_{H_n} + \big\|\xi^n_{1,i}\big\|_{H_n}.$$

Thus, for any $j\in I$, we deduce that $\big(\|\xi^n_{2,j}\|_{H_n}\big)_{n\geqslant 1}$ is also a bounded sequence.

Now, consider the ultraproduct $H=\prod_{\mathcal{U}} H_n$ of the Hilbert spaces H_n with respect to some ultrafilter $\mathcal{U}$ on $\mathbb{N}$ refining the Fréchet filter. For any $k\in\{1,2\}$ and any $i\in I$, let $\xi_{k,i}$ denote the vector of H corresponding to the sequence $\big(\xi^n_{k,i}\big)_{n\geqslant 1}$, i.e. the equivalence class of this sequence. The above computations give

$$\big\|\xi_{1,i}-\xi_{2,j}\big\|^2_{H} = a_{ij}, \quad i,j\in I. \tag{4.2}$$

For any $i,j\in I$ we let $\alpha_i=\xi_{1,i}$ and $\beta_j=\xi_{2,j}$. Then Equation (4.2) becomes

$$\big\|\alpha_i-\beta_j\big\|^2_{H} = a_{ij}, \quad i,j\in I.$$

Conversely, suppose that there exists a Hilbert space H and two families $(\alpha_i)_{i\in I}$ and $(\beta_j)_{j\in I}$ of elements of H such that for any $t\geqslant 0$ the Schur multiplier $T_t\colon B\big(\ell^2_I\big)\to B\big(\ell^2_I\big)$ is associated with the matrix

$$A_t = \Big[e^{-t\|\alpha_i-\beta_j\|^2_H}\Big]_{i,j\in I}.$$

Now, for any $t\geqslant 0$, we define the following matrices of $\mathbb{M}_I$

$$B_t = \Big[e^{-t\|\alpha_i-\alpha_j\|^2_H}\Big]_{i,j\in I},\quad C_t = \Big[e^{-t\|\beta_i-\beta_j\|^2_H}\Big]_{i,j\in I} \quad\text{and}\quad D_t = \Big[e^{-t\|\beta_i-\alpha_j\|^2_H}\Big]_{i,j\in I}.$$

For any $i\in I$ and any $n\in\{1,2\}$, we define the vector $\gamma_{(n,i)}$ of H by

$$\gamma_{(n,i)} = \begin{cases} \alpha_i & \text{if } n=1 \text{ and } i\in I \\ \beta_i & \text{if } n=2 \text{ and } i\in I. \end{cases}$$

Now, by the identification $\mathbb{M}_2(\mathbb{M}_I)\simeq\mathbb{M}_{\{1,2\}\times I}$, the block matrix $\begin{bmatrix} B_t & A_t \\ D_t & C_t \end{bmatrix}$ of $\mathbb{M}_2(\mathbb{M}_I)$ can be identified with the matrix

$$\Big[e^{-t\|\gamma_{(n,i)}-\gamma_{(m,j)}\|^2_H}\Big]_{(n,i)\in\{1,2\}\times I,(m,j)\in\{1,2\}\times I} \tag{4.3}$$

of $\mathbb{M}_{\{1,2\}\times I}$. Using [Arh1, Proposition 5.4], we deduce that, for any $t \geqslant 0$, the Schur multiplier associated with the matrix (4.3) is contractive on $B\big(\ell^2_{\{1,2\}\times I}\big)$. We deduce that $T_t\colon B\big(\ell^2_I\big) \to B\big(\ell^2_I\big)$ is also contractive. An alternative proof of this implication can be obtained in adapting the proof of [JMX, Proposition 8.17].

Finally, it is easy to see that $(T_t)_{t\geqslant 0}$ is a w*-continuous semigroup. ∎

Acknowledgment. The author would like to thank to Christian Le Merdy to provide him with the preprint [Xu] and Marius Junge and Eric Ricard for some discussions.

References

[Arh1] C. Arhancet. On Matsaev's conjecture for contractions on noncommutative L^p-spaces. Journal of Operator Theory 69, no. 2, 387–421 (2013).

[Arh2] C. Arhancet. Analytic semigroups on vector valued noncommutative L^p-spaces. Studia Math. 216 (2013), no. 3, 271–290.

[BCR] C. Berg, J. Christensen and P. Ressel. *Harmonic analysis on semigroups. Theory of positive definite and related functions*. Springer-Verlag, New York, 1984.

[Beu] A. Beurling. On analytic extension of semigroups of operators. J. Functional Analysis 6 (1970), 387–400.

[DCH] J. De Cannière and U. Haagerup. Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups. Amer. J. Math. 107, no. 2, 455–500, 1985.

[DJT] J. Diestel, H. Jarchow, and A. Tonge. Absolutely summing operators. Cambridge Studies in Advanced Mathematics 43, Cambridge University Press, Cambridge, 1995.

[EN] K.-J. Engel and R. Nagel. One-parameter semigroups for linear evolution equations. Graduate Texts in Mathematics, 194. Springer-Verlag, New York, 2000.

[ER] E. Effros and Z.-J. Ruan. Operator spaces. Oxford University Press (2000).

[Fac] S. Fackler. Regularity of semigroups via the asymptotic behaviour at zero. Semigroup Forum 87 (2013), no. 1, 117.

[Haa] M. Haase. The functional calculus for sectorial operators. Operator Theory: Advances and Applications, 169. Birkhäuser Verlag (2006).

[Hin] A. Hinrichs. K-convex operators and Walsh type norms. Math. Nachr. 208 (1999), 121–140.

[JMX] M. Junge, C. Le Merdy and Q. Xu. H^∞ functional calculus and square functions on noncommutative L^p-spaces. Astérisque No. 305 (2006).

[JuP] M. Junge and J. Parcet. The norm of sums of independent noncommutative random variables in $L_p(l_1)$. J. Funct. Anal. 221 (2005), no. 2, 366–406.

[NR] S. Neuwirth and É. Ricard. Transfer of Fourier multipliers into Schur multipliers and sumsets in a discrete group. Canad. J. Math. 63, no. 5, 1161–1187, 2011.

[Pau] V. Paulsen. Completely bounded maps and operator algebras. Cambridge Univ. Press (2002).

[Pis1] G. Pisier. Semi-groupes holomorphes et K-convexité. (French) [Holomorphic semigroups and K-convexity]. Seminar on Functional Analysis, 1980–1981, Exp. No. II, 32 pp., École Polytech., Palaiseau, 1981.

[Pis2] G. Pisier. Semi-groupes holomorphes et K-convexité (suite). (French) [Holomorphic semigroups and K-convexity (continued)] Seminar on Functional Analysis, 1980–1981, Exp. No. VII, 10 pp., École Polytech., Palaiseau, 1981.

[Pis3] G. Pisier. Holomorphic semigroups and the geometry of Banach spaces. Ann. of Math. (2) 115 (1982), no. 2, 375–392.

[Pis4] G. Pisier. Regular operators between non-commutative L_p-spaces. Bull. Sci. Math. 119 (1995), no. 2, 95–118.

[Pis5] G. Pisier. Non-commutative vector valued L_p-spaces and completely p-summing maps. Astérisque, 247 (1998).

[Pis6] G. Pisier. *Similarity problems and completely bounded maps*, volume 1618 of *Lecture Notes in Mathematics*. Springer-Verlag, expanded edition, 2001.

[Pis7] G. Pisier. Introduction to operator space theory. Cambridge University Press, Cambridge (2003).

[PX] G. Pisier and Q. Xu. Non-commutative L^p-spaces. 1459–1517 in *Handbook of the Geometry of Banach Spaces*, Vol. II, edited by W.B. Johnson and J. Lindenstrauss, Elsevier (2003).

[Ric] É. Ricard. A Markov dilation for self-adjoint Schur multipliers. Proc. Amer. Math. Soc. 136 (2008), no. 12, 4365–4372.

[SS] A. Sinclair and R. Smith. Finite von Neumann algebras and masas. London Mathematical Society Lecture Note Series, 351. Cambridge University Press, Cambridge, 2008.

[Xu] Q. Xu. H^∞ functional calculus and maximal inequalities for semigroups of contractions on vector-valued L_p-spaces. Preprint, arXiv:1402.2344.

Laboratoire de Mathématiques, Université de Franche-Comté, 25030 Besançon Cedex, France
cedric.arhancet@univ-fcomte.fr